

RESEARCH ARTICLE

The co-occurrence of *Callinectes sapidus* Rathbun, 1896 (Brachyura: Portunidae) and the parasitic dinoflagellate *Hematodinium* sp. (Dinoflagellata: Syndinidae) in two transitional water ecosystems of the Apulia coastline (South-Italy).

G. Mancinelli, L. Carrozzo, G. Marini*, P. Pagliara, M. Pinna

Department of Biological and Environmental Sciences and Technologies,
University of the Salento, S.P. Lecce-Monteroni, 73100 Lecce, Italy.

*Corresponding author: Phone: +39 0832 298604; E-mail address: gabriele.marini@unisalento.it

Abstract

- 1 - Coastal marine and transitional water ecosystems are threatened world-wide by invasions of non-indigenous invertebrate species, altering community structure and ecosystem functioning.
- 2 - Here, we report on the occurrence of the Atlantic blue crab *Callinectes sapidus* in transitional water ecosystems of the Apulia coastline (Salento peninsula, SE Italy).
- 3 - Blue crabs were sampled seasonally, every three months over a whole-year period in the Torre Colimena basin (Ionian Sea) and Acquatina lagoon (Adriatic Sea). Specimens were identified and enumerated. In addition, their hemolymph was screened for parasite infections using histological techniques.
- 4 - *Callinectes sapidus* was sampled in the Torre Colimena basin, where no other crab species were collected during the whole experimental period. In contrast, in the Acquatina Lagoon four crab species were sampled, including *C. sapidus*.
- 5 - The parasitic dinoflagellate *Hematodinium* sp., known to determine in *C. sapidus* the degenerative “bitter crab syndrome” in its native habitats, was ubiquitously detected in the hemolymph of blue crabs from both habitats and in autochthonous brachyurans collected in the Acquatina Lagoon.
- 6 - An assessment of the diverse spectrum of potential ecological effects triggered by the *Callinectes-Hematodinium* interaction on autochthonous brachyuran species is presented, together with a preliminary evaluation of the molecular approaches to be implemented in order to verify the species-specificity of the host-parasite interaction.

Keywords: *Callinectes sapidus*; not native species; *Hematodinium* sp.; bitter crab syndrome; host-parasite interaction; ecological impact; transitional water ecosystems; Salento peninsula.

Introduction

Originating from western Atlantic, the blue crab *Callinectes sapidus* Rathbun, 1896 (Decapoda, Brachyura, Portunidae) has to date extended its distribution in most European coastal waters and in Japan (Nehring, 2011). In Europe, the first record of *Callinectes* dates

back to 1900 on the Atlantic coast of France. Subsequently, specimens were detected in the North Sea (1932), Mediterranean Sea (1949, but probably as early as 1935), Baltic Sea (1951), Black Sea (1967), and possibly in the Sea of Azov (1967). In the Mediterranean Sea the species was first reported in Israel in

1955 (Holthuis and Gottlieb, 1955); in Italy it was observed in the Northern Adriatic Sea (Giordani Soika, 1951) and in the Genova Gulf (Tortonese, 1965). The majority of the reports on the blue crab in Mediterranean and, in general, European waters refers to episodic catches, limited in the number of specimens collected and the temporal period considered (Dulčić *et al.*, 2010; Nehring, 2011 and literature cited; but see Tureli Bilen *et al.*, 2011; Carrozzo *et al.*, 2013). In addition, virtually no information is available on the functional role played by this species within benthic food webs of Mediterranean coastal systems.

In its native habitats, *C. sapidus* represents an important component of estuarine communities (Baird and Ulanowicz, 1989; Dittel *et al.*, 2000). It has omnivorous, opportunistic trophic habits, feeding on plants, invertebrates, carcasses and conspecifics (Kennedy and Cronin, 2007); natural controls upon blue crab populations are represented by a number of large-sized fish predators including drums (e.g., *Pogonias cromis* and *Sciaenops ocellatus*), gars (e.g., *Lepisosteus spatula* and *L. oculatus*) and catfishes (e.g., *Ictalurus punctatus* and *I. furcatus*) (Darnell, 1961; Millikin and Williams, 1984). A further natural controlling agent is represented by the parasitic dinoflagellate *Hematodinium* sp., known to determine high mortalities in natural populations (Newman and Johnson, 1975; Messick and Shields, 2000; Pagenkopp Lohan *et al.*, 2012). In European waters, the pathogen has been described in a number of crustacean species, including brachyurans (Small *et al.*, 2012 and literature cited; Small, 2012), no information are available regarding its association with *C. sapidus* in the Mediterranean Sea yet.

In the present study, we surveyed the occurrence of *C. sapidus* in two coastal habitats located on the Adriatic and Ionian coasts of the Salento peninsula (Southern Italy). The survey was carried out over a whole

year using crab traps, allowing a previously unattempted quantitative estimation of seasonal variations in the species abundance in Mediterranean coastal habitats. In addition, we screened the hemolymph of sampled crab specimens in order to verify the occurrence of dinoflagellates belonging to the genus *Hematodinium* and the species-specificity of the parasite-host interaction.

Materials and Methods

Study sites

The study was carried out in two transitional water ecosystems of the South Apulia coastline (Salento Peninsula, SE Italy) i.e., the Acquatina lagoon and the Torre Colimena basin. The Acquatina lagoon (40°27'22"N, 18°12'24"E; Fig. 1) is a semi-artificial transitional water ecosystem created in the 1930s located on the Adriatic Sea. Information on the system morphology and oceanography is provided in Belmonte (2009), Mancinelli (2012), and Mancinelli *et al.* (2013). The lagoon is a shallow (1.2 m average depth) coastal basin with a surface area of 0.45 km² and about 2 km long. Dense stands of the seagrass *Cymodocea nodosa* (Ucria) Ascherson cover more than 50% of the bottoms, while other seagrasses of the genera *Ruppia* and *Zostera* together with macroalgae of the genera *Ulva*, *Ceramium*, *Chondria* and *Caulerpa* are locally abundant (Petrocelli and Cecere, 2009).

The benthic macrofauna and macroflora of the Acquatina lagoon have been extensively studied (De Mitri, 2004; Lumare *et al.*, 2009a, 2009b; Schirosi *et al.*, 2010; Mancinelli, 2012) and they are similar to other Adriatic transitional water ecosystems (Mancinelli and Rossi, 2001; Menéndez *et al.*, 2003; Mancinelli *et al.*, 2005; Mancinelli *et al.*, 2007a, 2007b; Orfanidis *et al.*, 2008; Ponti *et al.*, 2008; Basset *et al.*, 2012; Pinna *et al.*, 2013). Large-sized decapods are mainly represented by brachyuran species such as *Carcinus mediterraneus*, Czerniavsky




Figure 1. Location of the Torre Colimena basin and Acquatina lagoon in the Apulia coastline (SE Italy). Sampling stations are indicated by black circles and consecutive letters.

(Portunidae), *Eriphia verrucosa*, Forskal (Xantidae) and *Pachygrapsus marmoratus*, Fabricius (Grapsidae). *Maja verrucosa*, Milne-Edwards (Majidae) occurs episodically, while the hermit crab *Pagurus anachoretus*, Leach (Diogenidae) can be locally common. Detailed information on the Torre Colimena basin are provided in Carrozzo (2012) and in Carrozzo *et al.* (2013); in brief, the basin is a shallow (1.2 m average depth) artificial water body located along the Ionian coasts of the Salento Peninsula ($40^{\circ}17'59''\text{N}$; $17^{\circ}44'57''\text{E}$), with a surface area of 0.1 km² and about 0.35 km in diameter. It receives freshwater inputs from two drainage ditches,

while a channel connects the basin with the Ionian Sea (Fig. 1). The seagrass *C. nodosa* characterizes the soft bottom of the basin, while other seagrasses of the genera *Zostera* and *Ruppia* and macroalgae of the genera *Caulerpa* and *Chondria* are locally abundant. The benthic macrofauna is a brackish-water assemblage typical of other lagoons and coastal habitats of the Salento peninsula (e.g., Alemanno *et al.*, 2007a, 2007b, 2007c; Galuppo *et al.*, 2007; Mancinelli *et al.*, 2005, 2007a, 2008, 2009; Potenza and Mancinelli, 2010; Mancinelli, 2010; Vignes *et al.*, 2012). Among autochthonous brachyuran crustaceans, *C. mediterraneus* and *P.*

marmoratus are commonly observed along the banks of the basin.

Sampling procedures

Crabs were sampled with modified commercial crab traps (60 x 60 x 60 cm) of the type described by Sturdivant and Clark (2011). In brief, each trap was constructed of vinyl-coated 2 × 2 cm mesh wire with an upper and lower chamber. A 15 cm wide entrance was located on each of the four sides of the lower chamber, and a conical bait well was situated in the centre. The upper chamber was the parlour or trap chamber. The two chambers were separated by a wire-mesh panel, raised in the middle to form an inverted V. There were two openings along the apex of the V leading into the parlour. Each trap was attached to a line with a float for retrieval.

In April 2012, four sampling stations were located in each of the study habitats (Fig. 1). One crab trap was located at each sampling station at a depth of 70 - 100 cm. Each trap was baited with fish carcasses and fragmented mussels. Traps were deployed between 6.00 and 7.00 p.m. and retrieved the day after between 9.00 and 10.00 a.m. Crabs were harvested and transferred to the laboratory in refrigerated containers. Sampling operations were repeated adopting an identical procedure in summer (June), autumn (September), and winter (December).

Laboratory analyses

At each sampling time, captured crabs were identified and enumerated. Consequently, crab abundance was expressed in terms of catch per unit effort (CPUE hereafter) calculated as number of individuals per trap. Consequently, the hemolymph was extracted from each specimen from the arthrodistal membrane at the juncture of the basis and ischium of one of the swimming legs using a disposable 1ml syringe coupled to a 25-G

needle and containing an anti-coagulant solution (0.3 M NaCl, 0.1 M glucose, 30 mM sodium citrate, 26 mM citric acid, 10 mM EDTA, pH 5.4). One drop of hemolymph was mixed (1:1) with neutral red solution (0.25% w/v in filtered lagoon water) and examined under a light microscope. Parasites cells were stained by the solution and appeared red while host hemocytes remained translucent.

Results

*Occurrence of *Callinectes sapidus**

In total, 70 *C. sapidus* specimens were collected during the study period; 53 of these were captured in the Torre Colimena basin. In this habitat, no specimens belonging to other crab species were collected, even though during sampling operations both *P. marmoratus* Fabricius and *Carcinus* sp. were observed along the banks in crevices or other refuges. The blue crab was collected throughout the whole study period (Fig. 2); noticeably, peak abundance was observed in summer, while the occurrence of the species decreased considerably in winter and spring seasons.

In contrast, the 17 specimens of *C. sapidus* captured in the Acquatina Lagoon were sampled only in summer and autumn (Fig. 2), with maximal abundance approximately 50% lower than those observed in the Torre Colimena basin. A total of 20 specimens belonging to three other autochthonous brachyuran species were captured in crab traps during the study period at Acquatina lagoon, i.e., *E. verrucosa*, *P. marmoratus* and *C. mediterraneus*. Specifically, *E. verrucosa* was captured in summer and in autumn together with *C. sapidus*, while *P. marmoratus* and *C. mediterraneus* occurred in winter and spring (Fig. 2).

*Occurrence of the parasite *Hematodinium* sp.*

Overall, 70.4% of the crab specimens




Figure 2. Seasonal patterns of variation in the abundance (expressed as CPUE, catch per unit effort: Individuals Trap⁻¹) of *Callinectes sapidus* in the Torre Colimena basin and Acquatina lagoon during the study period.

examined were infected by the parasitic dinoflagellate *Hematodinium* sp. (Fig. 3). *Callinectes sapidus* resulted more infected than autochthonous crab species (71.6% vs. 57.1%) even though the difference was not statistically significant (Difference test for proportions, $P > 0.05$). Blue crabs from Acquatina Lagoon showed a lower degree of infection than conspecifics from Torre Colimena basin (74.1% vs. 62.5%), in turn more infected than autochthonous brachyurans (57.1%).

Discussion

The abundance of the blue crab *C. sapidus* was characterized in the two habitats under analysis by different seasonal patterns: while in the Acquatina lagoon it occurred exclusively in summer and autumn months,

in the Torre Colimena basin it was captured in crab traps throughout the year. A wide spectrum of biotic and abiotic factors may have determined the observed inter-site differences in the occurrence of *C. sapidus*; among others, temperature may have played a crucial role, since the Torre Colimena basin compared to the Acquatina lagoon was characterized during the study by warmer mean (23.35 ± 1.15 °C vs. 21.74 ± 1.08 °C, respectively; mean \pm SE) and winter water temperatures (12.21 ± 0.56 °C vs. 9.25 ± 0.41 °C, respectively; mean \pm SE). In its native habitats, *C. sapidus* is characterized by a period of quiescence in the cold season, overwintering buried in bottom sediments (Millikin and Williams, 1984; Kennedy and Cronin, 2007). The low water temperatures characterizing the Acquatina Lagoon and, in



Figure 3. Light micrographs of *Hematodinium* spp. infection in *Callinectes sapidus* hemolymph. Parasites (uninucleate trophonts) are stained by neutral red. Scale bar = 20 μm.

general, the southern Adriatic Sea may have replicated these environmental conditions; in contrast, the warmer Ionian waters are likely to have allowed the blue crab to extend its period of activity, bypassing the winter quiescence.

The abundance patterns observed for other autochthonous brachyurans indirectly suggest that negative interactions may occur between *C. sapidus* and native crab species, in particular those characterized by smaller sizes such as *P. marmoratus* and *C. mediterraneus*. The large-sized *Eriphia verrucosa*, conversely, seems to be less affected by the occurrence of the blue crab in the Acquatina Lagoon. In addition, it is apparent that in the Torre Colimena basin - characterized by a regular occurrence of the blue crab at high abundances - autochthonous crab species, even though present in

the benthic environment as testified by qualitative observations, did not occur in crab traps.

Indeed, strong interspecific competition has been observed between native and non-indigenous crab species in western Atlantic and eastern Pacific shore systems (McDonald *et al.*, 2001; Jensen *et al.*, 2002). Here, body size seems to influence the occurrence and magnitude of the interspecific interaction directly, or mediated by differential response related to species-specific variations in metabolism and, in turn, sensitivity to temperature fluctuations in the benthic environment. Further field and laboratory studies are needed to assess the nature and strength of the interactions (competition for trophic resources, or intra guild predation) linking the blue crab to other native brachyurans or, alternatively, the

influence of abiotic constraints in regulating the abundance and co-occurrence of crab species.

The second and most apparent finding of the present study is represented by the high frequency of the infection by the parasite *Hematodinium* sp. in the blue crab hemolymph, independently from the habitat, and the occurrence of the parasite in autochthonous brachyurans. Species of the genus *Hematodinium* are fast reproducing parasitic alveolates belonging to the class Syndinea. Within weeks from the first signs of infection by *Hematodinium*, the hemolymph of host crustaceans turns from opalescent to milky-white as parasite cells fill the hemal sinuses and replace crustacean hemocytes; muscle tissues of infected animals are degraded by the parasite, the hosts metabolic functions are severely compromised and death occurs within weeks of infection (“bitter crab syndrome”, BCS hereafter; Field *et al.*, 1992; Stentiford and Shields, 2005). In the United States and Canada, the BCS has raised considerable concern for commercially valuable crustacean species (Stentiford and Shields, 2005; Shields, 2012; Stentiford *et al.*, 2012b), as epizootics of *Hematodinium* spp. have severely affected a number of important fisheries. For example, in Alaskan tanner crabs (*Chionoecetes bairdi*) the parasite occurred in a third of fished districts, causing substantial decline in crab populations (Meyers *et al.*, 1990); similarly, the BCS has been recognized as one of the causes of the dramatic decline of the snow crab *Chionoecetes opilio* (Taylor and Khan, 1995; Pestal *et al.*, 2003). Most importantly, the *C. sapidus* fishery has suffered significant annual mortalities in high-salinity waters of Maryland and Virginia (Messick, 1994; Messick and Shields, 2000).

In European waters, infections have been observed in Scotland, where populations of *Nephrops norvegicus* have experienced *Hematodinium* infection rates of up to 70%

(Field *et al.*, 1992). In addition, the French velvet crab *Necora puber* fishery suffered a catastrophic reduction due to *H. perezii* (Wilhelm and Mialhe, 1996), and the edible crab *Cancer pagurus* fishery was also affected by *Hematodinium* spp. pathogens (Ní Chualáin *et al.*, 2009). *Hematodinium* parasites have been observed also in the caridean *Crangon crangon* (Stentiford *et al.*, 2012a).

In the Mediterranean Sea the parasite has been reported only episodically (Stentiford and Shields, 2005; Small *et al.*, 2012) while no information are to date available for Italian waters beside the present study. Here, the presence of *Hematodinium* in the hemolymph of *C. sapidus* and autochthonous brachyurans such as *C. mediterraneus*, *E. verrucosa* and *P. marmoratus* clearly indicates that the parasite may be a host generalist, a point elaborated in recent studies (Eigemann *et al.*, 2010; Pagenkopp Lohan *et al.*, 2012).

The importance of parasites in regulating animal invasions is widely acknowledged (Prenter *et al.*, 2004; Hatcher *et al.*, 2006; Wood *et al.*, 2007; Hatcher *et al.*, 2008; Lefèvre *et al.*, 2009; Dick *et al.*, 2010), and a huge spectrum of alternative hypotheses can be considered regarding the respective roles played by the blue crab, the native brachyuran species and the pathogen as preferential hosts, reservoir hosts and transmission routes. These alternative scenarios are briefly summarized in figure 4, ranging from apparent competition (A) to intra-guild predation (G). The information available to date do not allow to discriminate unambiguously among them; further efforts are needed to verify whether different species of *Hematodinium* occur among crab species, and the species-specificity of the infection. Only the identification of genetic similarities using PCR-based approaches and the construction of phylogenies of parasites from different host species can recognize potential alternate hosts and transmission




Figure 4. Possible interactions involving parasites: (A) apparent competition; (B) interspecific resource competition with shared parasite (including interference competition as denoted by horizontal arrows); (C) interspecific resource competition and a specialist (not shared) parasite; (D) predator–prey interaction with infected prey (the parasite and predator compete for prey resource); (E) predator–prey interaction with infected predator (conforming to a simple food chain); (F) predator and prey with shared parasite (as may occur for intermediate/definitive host/vector transmission relationships); (G) parasite infecting two intra-guild predators (with *Callinectes* preying more efficiently on native brachyurans). Modified from Hatcher *et al.* (2006).

routes (Archie *et al.*, 2009; Nagle *et al.*, 2009), ultimately identifying the ecological consequences of the parasite–host interaction in terms of spill-over and spill-back effects (Kelly *et al.*, 2009; Pagenkopp Lohan *et al.*, 2012).

In conclusion, this study presents novel quantitative information on the co-occurrence of the blue crab *C. sapidus* and its parasite dinoflagellate *Hematodinium* sp. In addition, it suggests an approach to the estimation of the ecological impact of alien species in Mediterranean waters, articulated on the integration of data on the spatial-temporal occurrence of the alien species itself together with information on the host–parasite interactions in which is involved. Ranging from a population to a cellular and molecular scale, the proposed approach has the potential to provide useful information on the mechanisms driving the

dispersal and establishment of the blue crab in the Mediterranean, on the effects exerted on autochthonous brachyuran species and, in general, on the biodiversity of benthic systems.

Acknowledgements

This study was performed in partial fulfillment of the requirements for a Master's Degree thesis in Community Ecology by LC. The authors wish to thank the Consorzio Speciale per la Bonifica Arneo for the support provided. Funding from FUR 2011–2012 to GM, MP and PP are acknowledged. This paper is dedicated to Sofia Mancinelli, thy eternal summer shall not fade.

References

- Alemanno S, Mancinelli G, Basset A 2007a. A patch use behaviour approach to model leaf litter breakdown in aquatic environment. *Transitional Waters Bulletin* 2: 47–57.
- Alemanno S, Mancinelli G, Basset A 2007b. Detritus processing in tri-trophic food chains:

- a modelling approach. *International Review of Hydrobiology* 92: 103-116.
- Alemanno S, Mancinelli G, Basset A 2007c. Effects of invertebrate patch use behaviour and detritus quality on reed leaf decomposition in aquatic systems: A modelling approach. *Ecological Modelling* 205: 492-506.
- Archie EA, Luikart G, Ezenwa VO 2009. Infecting epidemiology with genetics: a new frontier in disease ecology. *Trends in Ecology and Evolution* 24: 21-30.
- Baird D, Ulanowicz RE 1989. The seasonal dynamics of the Chesapeake Bay ecosystem. *Ecological Monographs* 59: 329-364.
- Basset A, Barbone E, Borja A, Brucet S, Pinna M, Quintana XD, Reizopoulou S, Rosati I, Simboura N 2012. A benthic macroinvertebrate size spectra index for implementing the Water Framework Directive in coastal lagoons in Mediterranean and Black Sea ecoregions. *Ecological Indicators*, 12: 72-83.
- Belmonte G 2009. Il lago di Acquatina: storia degli studi. *Thalassia Salentina* 31: 3-7.
- Carrozzo L 2012. The Blue Crab *Callinectes sapidus* in the Salento Peninsula (SE Italy): Occurrence, Trophic Position, Ecological Impact of an Alien Species in the Mediterranean Sea. LAP Lambert Academic Publishing, Germany.
- Carrozzo L, Carlino P, Costantini ML, Rossi L, Mancinelli G 2013. Seasonal abundance and trophic position of the Atlantic blue crab *Callinectes sapidus* Rathbun 1896 in a Mediterranean coastal habitat. *Journal of Sea Research*, in review.
- Darnell RM 1961. Trophic spectrum of an estuarine community, based on studies of Lake Pontchartrain, Louisiana. *Ecology* 42: 553-568.
- De Mitri R 2004. Pesci e crostacei decapodi del bacino di Acquatina (Lecce). *Thalassia Salentina* 27: 21-32.
- Dick JTA, Armstrong M, Clarke HC, Farnsworth KD, Hatcher MJ, Ennis M, Kelly A, Dunn AM 2010. Parasitism may enhance rather than reduce the predatory impact of an invader. *Biology Letters* 6: 636-638.
- Dittel AI, Epifanio CE, Schwalm SM, Fantle MS, Fogel ML 2000. Carbon and nitrogen sources for juvenile blue crabs *Callinectes sapidus* in coastal wetlands. *Marine Ecology Progress Series* 194: 103-112.
- Dulčić J, Dragičević B, Lipej L 2010. New record of the Blue Crab, *Callinectes sapidus* Rathbun, 1896, (Decapoda: Brachyura) in the Adriatic Sea. *Annales, Series Historia Naturalis* 20: 23-28.
- Eigemann F, Burmeister A, Skovgaard A 2010. *Hematodinium* sp. (Alveolata, Syndinea) detected in marine decapod crustaceans from waters of Denmark and Greenland. *Diseases of Aquatic Organisms* 92: 59-68.
- Field RH, Chapman CJ, Taylor AC, Neil DM, Vickerman K 1992. Infection of the Norway lobster *Nephrops norvegicus* by a *Hematodinium*-like species of dinoflagellate on the west coast of Scotland. *Diseases of Aquatic Organisms* 13: 1-15.
- Galuppo N, Maci S, Pinna M, Basset A 2007. Habitat types and distribution of benthic macroinvertebrates in a transitional water ecosystem: Alimini Grande (Puglia, Italy). *Transitional Waters Bulletin* 1(4): 9-19.
- Giordani Soika A 1951. Il *Neptunus pelagicus* (L.) nell'alto Adriatico. *Natura* 42: 18-20.
- Hatcher MJ, Dick JTA, Dunn AM 2006. How parasites affect interactions between competitors and predators. *Ecology Letters* 9: 1253-1271.
- Hatcher MJ, Dick JTA, Dunn AM 2008. A keystone effect for parasites in intraguild predation? *Biology Letters* 4: 534-537.
- Holthuis LB, Gottlieb E 1955. The occurrence of the american Blue Crab, *Callinectes sapidus* Rathbun, in Israel waters. *Bulletin of the Research Council of Israel* 5B: 154-156.
- Jensen GC, McDonald PS, Armstrong AD 2002. East meets west: competitive interactions between green crab *Carcinus maenas*, and native and introduced shore crab *Hemigrapsus* spp. *Marine Ecology Progress Series* 225: 251-262.
- Kelly DW, Paterson RA, Townsend CR, Poulin R, Tompkins DM 2009. Parasite spillback: a neglected concept in invasion ecology? *Ecology* 90: 2047-2056.
- Kennedy VS, Cronin LE 2007. *The Blue Crab: Callinectes sapidus*. Maryland Sea Grant College, College Park, Maryland.
- Lefèvre T, Lebarbenchon C, Gauthier-Clerc M, Missé D, Poulin R, Thomas F 2009. The ecological significance of manipulative parasites. *Trends in Ecology & Evolution* 24: 41-48.
- Lumare D, Lumare L, Scirocco T, Florio M, Lumare F 2009a. Composizione strutturale e dinamica del pescato nel lago di Acquatina. *Thalassia Salentina* 31: 63-82.
- Lumare L, Lumare D, Florio M, Scirocco T, Lumare F 2009b. I crostacei decapodi del lago

- di Acquatina: struttura delle popolazioni e ciclo riproduttivo. *Thalassia Salentina* 31: 83-102.
- Mancinelli G 2010. Intraspecific, size-dependent variation in the movement behaviour of a brackish-water isopod: a resource-free laboratory experiment. *Marine and Freshwater Behaviour and Physiology* 43: 321-337.
- Mancinelli G 2012. To bite, or not to bite? A quantitative comparison of foraging strategies among three brackish crustaceans feeding on leaf litters. *Estuarine, Coastal and Shelf Science* 110: 125-133.
- Mancinelli G, Rossi L 2001. Indirect, size-dependent effects of crustacean mesograzers on the Rhodophyta *Gracilaria verrucosa* (Hudson) Papenfuss: evidence from a short-term study in the Lesina Lagoon (Italy). *Marine Biology* 138: 1163-1173.
- Mancinelli G, Sabetta L, Basset A 2005. Short-term patch dynamics of macroinvertebrate colonization on decaying reed detritus in a Mediterranean lagoon (Lake Alimini Grande, Apulia, SE Italy). *Marine Biology* 148: 271-283.
- Mancinelli G, Sabetta L, Basset A 2007a. Colonization of ephemeral detrital patches by vagile macroinvertebrates in a brackish lake: a body size-related process? *Oecologia* 151: 292-302.
- Mancinelli G, Sabetta L, Basset A 2007b. On the influence of temporal resolution in mesh bag decomposition studies. *International Review of Hydrobiology* 92: 135-145.
- Mancinelli G, Pinna M, Basset A 2008. Spatio-temporal variability of macrozoobenthos size structure of a coastal lagoon: the influence of spectrum resolution. *Transitional Waters Bulletin* 2(2): 81-92.
- Mancinelli G, Vignes F, Sangiorgio F, Mastrolia A, Basset A 2009. On the potential contribution of microfungi to the decomposition of reed leaf detritus in a coastal lagoon: a laboratory and field experiment. *International Review of Hydrobiology* 94: 419-435.
- Mancinelli G, Vizzini S, Mazzola A, Maci S, Basset A 2013. Cross-validation of $\delta^{15}\text{N}$ and FishBase estimates of fish trophic position in a Mediterranean lagoon: the importance of the isotopic baseline. *Estuarine Coastal and Shelf Science*, in press.
- McDonald PS, Jensen GC, Armstrong DA 2001. The competitive and predatory impacts of the nonindigenous crab *Carcinus maenas* (L.) on early benthic phase Dungeness crab *Cancer magister* Dana. *Journal of Experimental Marine Biology and Ecology* 258: 39-54.
- Menéndez M, Carlucci D, Pinna M, Comin FA, Basset A 2003. Effect of nutrients on decomposition of *Ruppia cirrhosa* in a shallow coastal lagoon. *Hydrobiologia* 506-509: 729-735.
- Messick GA 1994. Hematodinium perezii infections in adult and juvenile blue crabs *Callinectes sapidus* from coastal bays of Maryland and Virginia, USA. *Diseases of Aquatic Organisms* 19: 77-82.
- Messick GA, Shields JD 2000. Epizootiology of the parasitic dinoflagellate *Hematodinium* sp. in the American blue crab *Callinectes sapidus*. *Diseases of Aquatic Organisms* 43: 139-152.
- Meyers TR, Botelho C, Koeneman TM, Short S, Imamura K 1990. Distribution of bitter crab dinoflagellate syndrome in southeast Alaskan tanner crabs, *Chionoecetes bairdi*. *Diseases of Aquatic Organisms* 9: 37-43.
- Millikin MR, Williams AB 1984. Synopsis of biological data on blue crab, *Callinectes sapidus* Rathbun. FAO Fisheries Synopsis 138, NOAA Technical Report, NMFS.
- Nagle L, Place AR, Schott EJ, Jagus R, Messick G, Pitula JS 2009. Real-time PCR-based assay for quantitative detection of *Hematodinium* sp. in the blue crab *Callinectes sapidus*. *Diseases of Aquatic Organisms* 84: 79-87.
- Nehring S 2011. Invasion History and Success of the American Blue Crab *Callinectes sapidus* in European and Adjacent Waters. In Galil BS, Clark PF, Carlton JT (eds) *In the Wrong Place - Alien Marine Crustaceans: Distribution, Biology and Impacts*. Springer, Netherlands, 607-624.
- Newman MW, Johnson CA 1975. A disease of blue crabs (*Callinectes sapidus*) caused by a parasitic dinoflagellate, *Hematodinium* sp. *The Journal of Parasitology* 61: 554-557.
- Ní Chualáin C, Hayes M, Allen B, Robinson M 2009. *Hematodinium* sp. in Irish *Cancer pagurus* fisheries: infection intensity as a potential fisheries management tool. *Diseases of Aquatic Organisms* 83: 59-66.
- Orfanidis S, Pinna M, Sabetta L, Stamatis N, Nakou K 2008. Variation of structural and functional metrics in macrophyte communities within two habitats of eastern Mediterranean coastal lagoons: natural versus human effects. *Aquatic Conservation: Marine and Freshwater Ecosystems* 18: S45-S61.

- Pagenkopp Lohan KM, Reece KS, Miller TL, Wheeler KN, Small HJ, Shields JD 2012. The role of alternate hosts in the ecology and life history of *Hematodinium* sp., a parasitic dinoflagellate of the blue crab (*Callinectes sapidus*). *Journal of Parasitology* 98: 73-84.
- Pestal GP, Taylor DM, Hoenig JM, Shields JD, Pickavance R 2003. Monitoring the prevalence of the parasitic dinoflagellate *Hematodinium* sp. in snow crabs *Chionoecetes opilio* from Conception Bay, Newfoundland. *Diseases of Aquatic Organism* 53: 67-75.
- Petrocelli A, Cecere E 2009. Il macrofitobenthos. *Thalassia Salentina* 31: 21-28.
- Pinna M, Marini G, Rosati I, Neto JM, Patricio J, Marques JC, Basset A 2013. The usefulness of large body-size macroinvertebrates in the rapid ecological assessment of Mediterranean lagoons. *Ecological Indicators* 29: 48-61.
- Ponti M, Pinna M, Basset A, Moncheva S, Trayanova A, Georgescu LP, Beqiraj S, Orfanidis S, Abbiati M 2008. Quality assessment of Mediterranean and Black Sea transitional waters: comparing responses of benthic biotic indices. *Aquatic Conservation: Marine and Freshwater Ecosystems* 18: S62-S75.
- Potenza L, Mancinelli G 2010. Body mass-related shift in movement behaviour in the isopod *Lekanesphaera hookeri* (Isopoda, Flabellifera): A laboratory study. *Italian Journal of Zoology* 77: 354-361.
- Prenter J, MacNeil C, Dick JTA, Dunn AM 2004. Roles of parasites in animal invasions. *Trends in Ecology & Evolution* 19: 385-390.
- Schiroso R, Musco L, Giangrande A 2010. Benthic assemblage of Acquafina Lake (South Adriatic Sea): present state and long-term faunistic changes. *Scientia Marina* 74: 235-246.
- Shields JD 2012. The impact of pathogens on exploited populations of decapod crustaceans. *Journal of Invertebrate Pathology* 110: 211-224.
- Small HJ 2012. Advances in our understanding of the global diversity and distribution of *Hematodinium* spp. - significant pathogens of commercially exploited crustaceans. *Journal of Invertebrate Pathology* 110: 234-246.
- Small HJ, Shields JD, Reece KS, Bateman K, Stentiford GD 2012. Morphological and molecular characterization of *Hematodinium perezii* (Dinophyceae: Syndiniales), a dinoflagellate parasite of the harbour crab, *Liocarcinus depurator*. *Journal of Eukaryotic Microbiology* 59: 54-66.
- Stentiford G, Bateman K, Small H, Pond M, Ungfors A 2012a. *Hematodinium* sp. and its bacteria-like endosymbiont in European brown shrimp (*Crangon crangon*). *Aquatic Biosystems* 8: 24.
- Stentiford GD, Neil DM, Peeler EJ, Shields JD, Small HJ, Flegel TW, Vlcek JM, Jones B, Morado F, Moss S, Lotz J, Bartholomay L, Behringer DC, Hauton C, Lightner DV 2012b. Disease will limit future food supply from the global crustacean fishery and aquaculture sectors. *Journal of Invertebrate Pathology* 110: 141-157.
- Stentiford GD, Shields JD 2005. A review of the parasitic dinoflagellates *Hematodinium* species and *Hematodinium*-like infections in marine crustaceans. *Diseases of Aquatic Organisms* 66: 47-70.
- Sturdivant SK, Clark KL 2011. An evaluation of the effects of blue crab (*Callinectes sapidus*) behavior on the efficacy of crab pots as a tool for estimating population abundance. *Fishery Bulletin* 109: 48-55.
- Taylor DM, Khan RA 1995. Observations on the occurrence of *Hematodinium* sp. (Dinoflagellata: Syndinidae), the causative agent of bitter crab disease in Newfoundland snow crab (*Chionoecetes opilio*). *Journal of Invertebrate Pathology* 65: 283-288.
- Tortonese E 1965. La comparsa di *Callinectes sapidus* Rath. (Decapoda, Brachyura) nel Mar Ligure. *Annali del Museo Civico di Storia Naturale di Genova* 4: 1-3.
- Tureli Bilen C, Korkcu P, Ibrikci T 2011. Application of artificial neural networks (ANNs) for weight predictions of blue crabs (*Callinectes sapidus* Rathbun, 1896) using predictor variables. *Mediterranean Marine Science* 12: 439-446.
- Vignes F, Fedele M, Pinna M, Mancinelli G, Basset A 2012. Variability of *Lekanesphaera monodi* metabolic rates with habitat trophic status. *Acta Oecologica* 41: 58-64.
- Wilhelm G, Mialhe E 1996. Dinoflagellate infection associated with the decline of *Necora puber* crab populations in France. *Diseases of Aquatic Organisms* 26: 213-219.
- Wood CL, Byers JE, Cottingham KL, Altman I, Donahue MJ, Blakeslee AMH 2007. Parasites alter community structure. *Proceedings of the National Academy of Sciences* 104: 9335-9339.